Economics: *see* Aphicide, 73
Economics: *see* Weed control, 337
Ecosystem: *see* Ground cover management, 65
Electrostatics: Deposition patterns and biological effectiveness of spray deposits on pot plants applied by the Ulvafan and three prototype electrostatic sprayers, 295
Euphorbia heterophylla: *see* Herbicide, 218
European red mite: *see Panonychus ulmi*, 463
Eyespot: *see Pseudocercosporella herpotrichoides*, 135, 265
Fenvalerate: *see Psylla pyricola*, 185
Fertilizer: *see* Cloves, 415
Field resistance: *see Diuraphis noxia*, 255
Flat-fan nozzle: *see* Spray volume distribution, 111
Flowering: *see Monolepta australis*, 392
Forecasting: Forecasting future developments in crop protection, 83
Forecasting: *see* Aphicide, 73
Forest weed species: *see* Phytotoxicity, 52
Formulations: *see* Ultraviolet light, 21
Fruit-set: *see Monolepta australis*, 392
Fungicidal control: *see Septoria apiicola*, 433
Fungicide resistance: *see* Benomyl, 190
Fungicide–herbicide interaction: Interactions of fungicide–herbicide combinations against plant pathogens and weeds, 403
Fungicide: *see* Cocoa, 281
Fungicide: *see* Leaf blotch, 367
Fungicide: *see Lycopersicon esculentum*, 231
Fungicide: *see* Narcissus, 143
Fungicide: *see Phytophthora infestans*, 15
Fungicide: *see Pseudocercosporella herpotrichoides*, 135, 265
Fungicide: *see Rhizoctonia* root rot, 275
Fungicide: *see* Seed treatment, 439

Gall midge: *see Orseolia oryzivora*, 252
Gliocladium virens: *see* Tomato fruit rot, 8
Glycine max: Method of assessment of red leaf blotch on soybeans used to evaluate cultivar susceptibility and chemical control, 148
Glyphosate: *see* Imazapyr, 59
Glyphosate: *see Oscinella* spp., 105
Glyphosate: *see Orobanche cernua*, 332
Glyphosate: *see* Phytotoxicity, 52
Gossypium hirsutum: *see Verticillium* wilt, 174
Grass reseed: *see Oscinella* spp., 105
Ground cover management: Effects of ground cover managment on certain abiotic and biotic interactions in peach orchard ecosystems, 65
Ground cover: *see Aphis pomi*, 225

Hardwood: *see* Imazapyr, 59
Helicoverpa armigera: *see* Chickpea, 29
Helicoverpa armigera: *see* Model, 197
Herbicide ring weeding: *see* Cloves, 415
Herbicide: Control of *Euphorbia heterophylla* L. in cowpea with herbicides and herbicide mixtures, 218
Herbicide: *see Lolium rigidum*, 422
Herbicide: *see Orobanche cernua*, 332
Herbicide: *see s*-triazine, 429
Herbicide: *see* Weed threshold, 323
Hexythiazox: *see Panonychus ulmi*, 463
Hymexazol: Hymexazol treatment of sugar-beet seed to control seedling disease caused by *Pythium* spp. and *Aphanomyces cochlioides*, 371

Imazapyr: Stump treatment with imazapyr and glyphosate after cutting hardwoods to prevent regrowth of suckers, 59
Imazapyr: *see* Phytotoxicity, 52
Insect control: *see* Model, 197
Insect resistance: *see Nicotiana tabacum*, 351
Insecticidal control: *see Monolepta australis*, 392
Insecticide residues: Some factors influencing the accelerated degradation of mephosfolan in soils, 44
Insecticide resistance: *see Psylla pyricola*, 185
Insecticide: *see Aphis gossypii*, 39
Insecticide: *see* Chickpea, 29
Insecticide: *see Orseolia oryzivora*, 252
Insecticide: *see Psila rosae*, 395
Insecticide: *see Somaticus*, 128
Integrated control: *see Lycopersicon esculentum*, 231
Integrated control: *see Verticillium dahliae*, 363
Integrated management: *see* Citrus bacterial canker disease, 3
Integrated pest management: *see* Noctuid larvae, 115
Integrated pest management: *see Oscinella frit*, 97
Irritancy: *see Aphis gossypii*, 39
Isazofos: *see Orseolia oryzivora*, 252
Isoproturon: Measurement and prediction of isoproturon movement and persistence in three soils, 289

Jembe ring weeding: *see* Cloves, 415
Juvenile decline: *see* Cloves, 415

Lactuca sativa: Screening for resistance to big-vein disease of lettuce (*Lactuca sativa*), 446
Leaching: *see* Isoproturon, 289
Leaf blotch: Tests of fungicides for the control of leaf blotch diseases: *Cladosporium allii* of leek and *C. alliicepae* of onion, 367
Lettuce big-vein virus: *see Lactuca sativa*, 446
Lolium multiflorum: *see Oscinella frit*, 101
Lolium rigidum: Model of wheat yield response to application of diclofop-methyl to control ryegrass (*Lolium rigidum*), 422
Lycopersicon esculentum: Integrated control of *Sclerotium rolfsii* infection of tomato in the Nigerian Savanna: effect of *Trichoderma viride* and some fungicides, 231

Maize: *see s*-triazine, 429
Maize: *see* Somaticus, 128
Management: *see Papaipema nebris*, 387
Meloidogyne spp.: *see* Nematodes, 453
Mephosfolan: *see* Insecticide residues, 44
Mirid head bug: Biology and population dynamics of sorghum head bugs (Hemiptera: Miridae), 164
Mites: *see* Stored product, 378
Miticides: *see Tetranychus urticae*, 119
Model: Use of simulation analysis to assess *Helicoverpa armigera* control on pigeonpea in southern India, 197
Model: *see* Weed control, 337
Model: *see* Weed threshold, 323
Modified atmosphere: *see* Biogenerators, 235
Monilinia fructicola: *see* Benomyl, 190
Monolepta australis: Invasion of flowering avocado trees by *Monolepta australis* (Jacoby) (Coleoptera: Chrysomelidae) and its impact on fruit-set in North Queensland, 392

Narcissus: Effect of fungicides on the production of adventitious bulbils in the propagation of *Narcissus* by the chipping technique, 143
Nematodes: Global status of nematode problems of groundnut, pigeonpea, chickpea, sorghum and pearl millet and suggestions for future work, 453
Nicotiana tabacum: Additive protective effects of different plant-derived insect resistance genes in transgenic tobacco plants, 351
Noctuid larvae: Evaluation of chemical and cultural control for noctuid larvae in Chilean artichokes produced for foreign markets, 115
Norway spruce: *see* Phytotoxicity, 52

Olpidium brassicae: *see Lactuca sativa*, 446
Orobanche cernua: Control of broomrape (*Orobanche cernua*) in sunflower (*Helianthus annuus* L.) with glyphosate, 332
Orseolia oryzivora: Efficacy of granular isazofos against rice gall midge in Nigeria, 252
Oryctolagus cuniculus: Effect of grazing by wild rabbits (*Oryctolagus cuniculus*) on the growth and yield of oilseed and fodder rape (*Brassica napus* sub. sp. oleifera), 155
Oryza sativa: Relative susceptibility of high-yielding rice varieties to the thrips, *Stenchaetothrips biformis* (Bagnall) (Thysanoptera: Thripidae), 193
Oryza sativa: *see Cnaphalocrocis medinalis*, 243
Oscinella frit: Control of frit fly (*Oscinella* spp.) in newly sown grass using a combination of low doses of pesticide, resistant grass cultivars and indigenous parasitoids, 97

Oscinella frit: Influence of date of sowing on frit-fly damage to newly sown Italian ryegrass, 101
Oscinella spp.: Origin of frit-fly (*Oscinella* spp.) attack on newly sown grass, effect of pesticides on frit-fly larvae and benefit of using glyphosate herbicide pre-ploughing, 105
Oxadixyl: *see Phytophthora infestans*, 15

Panolis flammea: *see* Ultraviolet light, 21
Panonychus ulmi: Comparison of clofentezine and hexytiazox with cyhexatin for integrated control of European red mite, *Panonychus ulmi* (Koch.), 463
Papaipema nebris: Evaluation of two management strategies for stalk borer, *Papaipema nebris*, in corn, 387
Parasitic weed: *see Orobanche cernua*, 332
Peach tree: *see* Ground cover management, 65
Pear psylla: *see Psylla pyricola*, 185
Permethrin: *see Papaipema nebris*, 387
Persistence: *see s-triazine*, 429
Pest damage: *see Oscinella frit*, 101
Phytophthora infestans: Competition between oxadixyl-sensitive and -resistant field isolates of *Phytophthora infestans* on fungicide-treated potato crops, 15
Phytotoxicity: Effect of foliage-applied imazapyr and glyphosate on common forest weed species and Norway spruce, 52
Pigeonpea: *see* Model, 197
Plant density: *see Verticillium* wilt, 174
Pod damage-yield relationship: *see* Chickpea, 29
Population dynamics: *see* Mirid head bug, 164
Pot plant: *see* Electrostatics, 295
Potato: *see Phytophthora infestans*, 15
Potato: *see Verticillium dahliae*, 363
Pratylenchus spp.: *see* Nematodes, 453
Propagation: *see Narcissus*, 143
Propiconazole: *see* Leaf blotch, 367
Pseudocercosporella herpotrichoides: Changes in populations of *Pseudocercosporella herpotrichoides* in successive crops of winter wheat in relation to initial populations and fungicide treatments, 135
Pseudocercosporella herpotrichoides: Effects of fungicide timing on the severity of eyespot on winter wheat or winter barley inoculated with W-type or R-type isolates of *Pseudocercosporella herpotrichoides*, 265
Pseudocercosporella herpotrichoides: *see* Fungicide–herbicide interaction, 403
Psila rosae: Control of carrot fly (*Psila rosae* F.) in celeriac, 395
Psylla pyricola: Resistance to the pyrethroid fenvalerate in pear psylla, *Psylla pyricola* Foerster (Homoptera: Psyllidae), in the North-western USA, 185
Pyrenochaeta glycines: *see Glycine max*, 148
Pythium spp.: *see* Hymexazol, 371

Rabbit: *see Oryctolagus cuniculus*, 155
Red leaf blotch: *see Glycine max*, 148
Redshouldered leaf beetle: *see Monolepta australis*, 392
Resistance breeding: *see Nicotiana tabacum*, 351
Resistance screening: *see Lactuca sativa*, 446
Resistance: *see* Citrus bacterial canker disease, 3
Resistance: *see Oryza sativa*, 193
Resistance: *see Phytophthora infestans*, 15
Rhizoctonia cerealis: *see* Fungicide–herbicide interaction, 403
Rhizoctonia root rot: Evaluation of methods for the chemical control of rhizoctonia root rot of wheat, 275
Rhizoctonia solani: *see* Compost, 271
Rhizoctonia solani: *see* Tomato fruit rot, 8
Rice leaffolder: see *Cnaphalocrocis medinalis*, 243
Rice varieties: *see Oryza sativa*, 193
Rice: *see Cnaphalocrocis medinalis*, 243
Rice: *see Orseolia oryzivora*, 252
Rosellinia necatrix: *see* Solarization, 312
Roses: *see Tetranychus urticae*, 119
Rotary atomization: Measurement of drop spectra from rotary cage aerial atomizers, 33
Russian wheat aphid: *see Diuraphis noxia*, 255
Ryegrass *see Oscinella frit*, 97

s-triazine: Triazine herbicide carry-over in semi-arid conditions, 429
Sclerotium rolfsii: *see Lycopersicon esculentum*, 231
Sclerotium rolfsii: *see* Compost, 271
Seed treatment: Improving wheat seedling emergence by seed-protectant fungicides, 439
Seed treatment: *see* Hymexazol, 371
Seedling emergence: *see* Seed treatment, 439
Semi-arid tropics: *see s*-triazine, 439
Septoria apiicola: Glasshouse and field evaluation of fungicides for the control of *Septoria apiicola* on celery, 433
Sinapis alba: *see* Fungicide–herbicide interaction, 403
Sitobion avenae: *see* Aphicide, 73
Slug control: Laboratory and field assessment of a new aluminium chelate slug poison, 131
Slug: Assessment of slug activity using bran-baited traps, 355
Soil: *see* Insecticide residues, 44
Soil: *see* Isoproturon, 289
Solarization: Long-term effect of soil solarization for the control of *Rosellinia necatrix* in apple, 312
Somaticus: Efficacy of soil-applied and seed dressing insecticides for the control of false wireworms, *Somaticus* species (Coleoptera: Tenebrionidae), in maize, 128
Sorghum: *see* Mirid head bug, 164
Sowing date: *see Oscinella frit*, 101
Soybean: *see Glycine max*, 148
Spatial distribution: *see* Weed control 337
Spodoptera exempta: Effects of azadirachtin and acqueous neem seed extracts on survival, growth and development of the African armyworm, *Spodoptera exempta*, 383
Spray contamination: Effect of application volume and method on spray operator contamination by insecticide during cotton spraying, 343
Spray deposit: *see* Electrostatics, 295
Spray volume distribution: Predicting the transverse volume distribution under an agricultural spraying boom, 111
Stalk borer: *see Papaipema nebris*, 387
Stenchaetothrip biformis: *see Oryza sativa*, 193
Stone fruit: *see* Benomyl, 190
Stored product: Preventive biological control of stored food mites in empty stores using *Cheyletus eruditus* (Schrank), 378
Stored product: *see* Biogenerators, 235
Sugar beet: *see* Hymexazol, 371
Sunflower: *see Agelaius phoeniceus*, 359
Sunflower: *see Orobanche cernua*, 332
Synergism: *see* Fungicide–herbicide interaction, 403

Tenacity: *see* Cocoa, 281
Tetranychus urticae: Evaluation of miticides for the control of two-spotted mite *Tetranychus urticae* Koch on field roses in southern Queensland, 119
Threshold: *see* Weed control, 337
Thrips tabaci: Investigation of *Thrips tabaci* Lind. as a cotton pest and the development of strategies for its control in the Punjab, 469
Thrips: *see Oryza sativa*, 193
Time-lapse video: *see* Slug, 355
Tomato fruit rot: Greenhouse and field studies on the biological control of tomato fruit rot caused by *Rhizoctonia solani*, 8
Transgenic plant: *see Nicotiana tabacum*, 351
Trap crop: Effect of mustard (*Brassica hirta*) as a non-crop plant on competition and insect pests in broccoli (*Brassica oleracea*), 90
Trap: *see* Slug, 355
Trichoderma hamatum: *see* Tomato fruit rot, 8
Trichoderma harzianum: *see Verticillium dahliae*, 363
Trichoderma viride: *see Lycopersicon esculentum*, 231
Two-spotted mite: *see Tetranychus urticae*, 119
Typhlodromus pyri: *see Panonychus ulmi*, 463

Ultra-low-volume spraying: *see* Application method, 207

Ultraviolet light: Influence of drop size, solar ultraviolet light and protectants, and other factors on the efficacy of baculovirus sprays against *Panolis flammea* (Schiff.) (Lepidoptera: Noctuidae), 21

Varietal disease: *see Cnaphalocrocis medinalis*, 243
Verticillium dahliae: Integrated control of *Verticillium dahliae* in potato by *Trichoderma harzianum* and captan, 363
Verticillium lecanii: Comparison of ultra-low-volume electrostatic and high-volume hydraulic application of *Verticillium lecanii* for aphid control in chrysanthemums, 177
Verticillium wilt: Inverse relationship between cotton plant density and verticillium wilt incidence and severity: evidence for an alternative hypothesis, 174

Weed control: Spatial weed distribution and economic thresholds for weed control, 337
Weed threshold: Weed economic thresholds in cereals in the Federal Republic of Germany, 323
Weed, pasture: *see* Allelopathy, 410
Wheat: *see Lolium rigidum*, 422
Wheat: *see* Rhizoctonia root rot, 275
Wheat: *see* Seed treatment, 439
Wheat; bread: *see Diuraphis noxia*, 255
Wheat, winter, *see* Aphicide, 73
Wheat, winter: *see Branta bernicla*, 453
Wheat, winter: *see Pseudocercosporella herpotrichoides*, 135, 265
White root rot: *see* Solarization, 312
Wild asparagus: *see Brachycorynella asparagi*, 259
Wind tunnel: *see* Rotary atomization, 33
Wireworm: *see Somaticus*, 128

Xanthomonas campestris: *see* Citrus bacterial canker disease, 3

Zea mays: *see Papaipema nebris*, 387

Author Index

ADAMS, A. J. AND HALL, F. R. Initial behavioural responses of *Aphis gossypii* to defined deposits of bifenthrin on chrysanthemum, 39
AHMAD, Z. *see* Attique, M. R., 469
AKINYEMIJU, O. A. *see* O'MAKINWA, R. O., 218
ALTIERI, M. A. *see* KLOEN, H., 90
ARAYA, J. E. *see* MACHUCA, J. R., 155
ARMES, N. J. *see* HOLT, J., 197
ARRETZ V. P. *see* MACHUCA, J. R., 155
ASTERAKI, E. J. *see* CLEMENTS, R. O, 105
ATTIQUE, M. R. AND AHMAD, Z. Investigation of *Thrips tabaci* Lind. as a cotton pest and the development of strategies for its control in the Punjab, 469
ATTIQUE, M. R. *see* SUTHERLAND, J. A., 343

BALLINGER, D. J. *see* COTTERILL, P. J., 275
BARKSDALE, T. H. *see* LEWIS, J. A., 8
BATEMAN, G. L., FITT, B. D. L., CREIGHTON, N. F. AND HOLLOMON, D. W. Changes in populations of *Pseudocercosporella herpotrichoides* in successive crops of winter wheat in relation to initial populations and fungicide treatments, 135
BENTLEY, B. R. *see* CLEMENTS, R. O., 101
BENTLEY, B. R. *see* CLEMENTS, R. O., 105
BERRIE, A. M. *see* CROSS, J. V., 207
BLAIR, A. M., MARTIN, T. D., WALKER, A. AND WELCH, S. J. Measurement and prediction of isoproturon movement and persistence in three soils, 289
BOAG, B., MACFARLANE SMITH, W. H. AND GRIFFITHS, D. W. Effect of grazing by wild rabbits (*Oryctolagus cuniculus*) on the growth and yield of oilseed and fodder rape (*Brassica napus* sub. sp. *oleifera*), 155
BOS, L. AND HUIJBERTS, N. Screening for resistance to big-vein disease of lettuce (*Lactuca sativa*), 446
BOULTER, D., EDWARDS, G. A., GATEHOUSE, A. M. R, GATEHOUSE, J. A. AND HILDER, V. A. Additive protective effects of different plant-derived insect resistance genes in transgenic tobacco plants, 351
BOWER, C. C. Comparison of clofentezine and hexythiazox with cyhexatin for integrated control of European red mite, *Panonychus ulmi* (Koch.), 463
BRODY, A. K., KARBAN, R. AND SCHNATHORST, W. C. Inverse relationship between cotton plant density and verticillium wilt incidence and severity: evidence for an alternative hypothesis, 174
BURCHILL, R. T. see JORDAN, M. M., 367
BURTS, E. C. *see* VAN DE BAAN, H. E., 185

CALDERON, M. *see* PASTER, N., 235
CARTER, N. *see* WRATTEN, S. D., 73
CASTEJON-MUÑOZ, M., ROMERO-MUÑOZ, F. AND GARCÍA-TORRES, L. Control of broomrape (*Orobanche cernua*) in sunflower (*Helianthus annuus* L.) with glyphosate, 332
CAUSSIN R. *see* LEUNDA, P., 111
CHET, I. *see* ORDENTLICH, A., 363
CHIMA WOKOCHA, R. Integrated control of *Sclerotium rolfsii* infection of tomato in the Nigerian Savanna: effect of *Trichoderma viride* and some fungicides, 231
CHIVINGE, O. A. AND MPOFU, B. Triazine herbicide carryover in semi-arid conditions, 429
CLEAR HILL, B. H., VAN EMDEN, H. F. AND CLEMENTS, R. O. Control of frit fly (*Oscinella* spp.) in newly sown grass using a combination of low doses of pesticide, resistant grass cultivars and indigenous parasitoids, 97
CLEMENTS, R. O. *see* CLEAR HILL, B. H., 97
CLEMENTS, R. O., BENTLEY, B. R. AND JACKSON, C. A. Influence of date of sowing on frit-fly damage to newly sown Italian ryegrass, 101
CLEMENTS, R. O., BENTLEY, B. R. AND ASTERAKI, E. J. Origin of frit-fly (*Oscinella* spp.) attack on newly sown grass, effect of pesticides on frit-fly larvae and benefit of using glyphosate herbicide pre-ploughing, 105
COHEN, Y. AND SAMOUCHA, Y. Competition betwen oxadixyl-sensitive and -resistant field isolates of *Phytophthora infestans* on fungicide-treated potato crops, 15
COLE, D. L. *see* LEVY, C., 148
COOKE, B. K., HISLOP, E. C., HERRINGTON, P. J., WESTERN N. M. AND HUMPHERSON-JONES, F. Air-assisted spraying of arable crops, in relation to deposition, drift and pesticide performance, 303
COTTERILL, P. J., BALLINGER, D. J. AND KOLLMORGEN, J. F. Evaluation of methods for the chemical control of rhizoctonia root rot of wheat, 275
CREIGHTON, N. F. *see* BATEMAN, G. L., 135
CROFT, B. A. *see* VAN DE BAAN, H. E., 185
CROSS, J. V. AND BERRIE, A. M. Efficacy of reduced volume and reduced dose rate spray programmes in apple orchards, 207

DAVIS, P. M. AND PEDIGO, L. P. Evaluation of two management strategies for

stalk borer, *Papaipema nebris*, in corn, 387
DEFAVERI, S. G. *see* FAY, H. A. C., 392
DEBOUCHE, C. *see* LEUNDA, P., 111
DENT, J. B. *see* THORNTON, P. K., 337
DOBSON, H. M. *see* SUTHERLAND, J. A., 343
DRINKWATER, T. W., GILIOMEE, J. H. AND PRINGLE, K. L. Efficacy of soil-applied and seed dressing insecticides for the control of false wireworms, *Somaticus* species (Coleoptera: Tenebrionidae), in maize, 128
DU TOIT, F. Field resistance in three bread wheat lines to the Russian wheat aphid, *Diuraphis noxia* (Hemiptera: Aphididae), 255

EDWARDS, G. A. *see* BOULTER, D., 351
ENTWISTLE, J. C. *see* WRATTEN, S. D., 73

FAWCETT, R. H. *see* THORNTON, P. K. 337
FAY, H. A. C. AND DEFAVERI, S. G. Invasion of flowering avocado trees by *Monolepta australis* (Jacoby) (Coleoptera: Chrysomelidae) and its impact on fruit-set in North Queensland, 392
FITT, B. D. L. *see* BATEMAN, G. L., 135
FITT, B. D. L. *see* GOULDS, A., 265
FOLWELL, R. J., GEFRE, J. A., LUTZ, S. M., HALFHILL, J. E. AND TAMAKI, G. Methods and costs of suppressing *Brachycorynella asparagi Mordvilko* (Homoptera: Aphididae), 259
FREEMAN, S., SZTEJNBERG, A., SHABI, E. AND KATAN, J. Long-term effect of soil solarization for the control of *Rosellinia necatrix* in apple, 312

GARCÍA-TORRES, L. *see* CASTEJON-MUÑOZ, M., 332
GATEHOUSE, A. M. R. *see* BOULTER, D., 351
GATEHOUSE, J. A. *see* BOULTER, D., 351
GEFRE, J. A. *see* FOLWELL, R. J., 259
GEROWITT, B. AND HEITEFUSS, R. Weed economic thresholds in cereals in the Federal Republic of Germany, 323
GILIOMEE, J. H. *see* DRINKWATER, T. W., 128
GILLESPIE, A. T. *see* SOPP, P. I., 177
GISI, U. *see* KATARIA, H. R., 403
GORODECKI, B. AND HADAR, Y. Suppression of *Rhizoctonia solani* and *Sclerotium rolfsii* diseases in container media containing composted separated cattle manure and composted grape marc, 271
GOTSCH, N. AND RIEDER, P. Forecasting future developments in crop protection, 83
GOUGH, N. Evaluation of miticides for the control of two-spotted mite *Teiranychus urticae* Koch on field roses in southern Queensland, 119
GOULDS, A. AND FITT, B. D. L. Effects of fungicide timing on the severity of eyespot on winter wheat or winter barley inoculated with W-type or R-type isolates of *Pseudocercosporella herpotrichoides*, 265
GRIFFITHS, D. W. *see* BOAG, B., 155

HADAR, Y. *see* GORODECKI, B., 271
HADLEY, P. *see* RUDGARD, S. A., 281
HALEY, S. AND HOGUE, E. J. Ground cover influence on apple aphid, *Aphis pomi* DeGeer (Homoptera: Aphididae) and its predators in a young apple orchard, 225
HALFHILL, J. E. *see* FOLWELL, R. J., 259
HALL, F. R. *see* ADAMS, A. J., 39
HANZEL, J. J. *see* MAH, J., 359
HEITEFUSS, R. *see* GEROWITT, B., 323
HENDERSON, I. F., MARTIN, A. P. AND PARKER, K. A. Laboratory and field assessment of a new aluminium chelate slug poison, 131
HERRINGTON, P. J. see COOKE, B. K., 303
HILDER, V. A. *see* BOULTER, D., 351
HILLMAN, G. *see* SUMMERS, R. W., 459
HISLOP, E. C. *see* COOKE, B. K., 303
HOGUE, E. J. *see* HALEY, S., 225
HOLLOMON, D. W. *see* BATEMAN, G. L., 135
HOLT, J., KING, A. B. S. AND ARMES, N. J. Use of simulation analysis to assess *Helicoverpa armigera* control on pigeonpea in southern India, 197
HORÁK, E. *see* ŽĎÁRKOVÁ, E., 378
HUIJBERTS, N. *see* BOS, L., 446
HUMPHERSON-JONES, F. *see* COOKE, B. K., 303

INGRAM, W. R. *see* SUTHERLAND, J. A., 343

JACKSON, C. A. *see* CLEMENTS, R. O., 101
JORDAN, M. M., MAUDE, R. B. AND BURCHILL, R. T. Tests of fungicides for the control of leaf blotch diseases: *Cladosporium allii* of leek and *C. alliicepae* of onion, 367
JOSHI, R. C. *see* KHAN, Z. R., 243
JUKES, A. A. *see* SUETT, D. L., 44

KARBAN, R. *see* BRODY, A. K., 174
KATAN, J. *see* FREEMAN, S., 312
KATARIA, H. R. AND GISI, U. Interactions of fungicide-herbicide combinations against plant pathogens and weeds, 403
KHAN, Z. R. AND JOSHI, R. C. Varietal resistance to *Cnaphalocrocis medinalis* (Guenee) in rice, 243
KILLICK, H. J. Influence of drop size, solar ultraviolet light and protectants, and other factors on the efficacy of baculovirus sprays against *Panolis flammea* (Schiff.) (Lepidoptera: Noctuidae), 21
KING, A. B. S. *see* HOLT, J., 197
KING, W. J. *see* SUTHERLAND, J. A., 343
KLOEN, H. AND ALTIERI, M. A. Effect of mustard (*Brassica hirta*) as a non-crop plant on competition and insect pests in broccoli (*Brassica oleracea*), 90
KOLLMORGEN, J. F. *see* COTTERILL, P. J., 275

LARRAIN, P. I. *see* MACHUCA, J. R., 115
LEITE, JR, R. P. AND MOHAN, S. K. Integrated management of the citrus bacterial canker disease caused by *Xanthomonas campestris* pv. *citri* in the State of Parana, Brazil, 3
LEUNDA, P., DEBOUCHE, C. AND CAUSSIN, R. Predicting the transverse volume distribution under an agricultural spraying boom, 111
LEVY, C., MAHUKU, G. S, TATTERSFIELD, J. R. AND COLE, D. L. Method of assessment of red leaf blotch on soybeans used to evaluate cultivar susceptibility and chemical control, 148
LEWIS, J. A., BARKSDALE, T. H. AND PAPAVIZAS, G. C. Greenhouse and field studies on the biological control of tomato fruit rot caused by *Rhizoctonia solani*, 8
LINFIELD, C. A. AND PRICE, D. Effect of fungicides on the production of adventitious bulbils in the propagation of Narcissus by the chipping technique, 143
LINZ, G. M. *see* MAH, J., 359
LISKER, N. Improving wheat seedling emergence by seed-protectant fungicides, 439
LOPEZ, V. F. *see* SHARMA, H. C., 164
LUND-HØIE, K. AND ROGNSTAD, A. Effect of foliage-applied imazapyr and glyphosate on common forest weed species and Norway spruce, 52
LUND-HØIE, K. AND ROGNSTAD, A. Stump treatment with imazapyr and glyphosate after cutting hardwoods to prevent regrowth of suckers, 59
LUTZ, S. M. *see* FOLWELL, R. J., 259

MACFARLANE SMITH, W.H. *see* BOAG, B., 155
MACHUCA, J. R., ARAYA, J. E., ARRETZ, V. P. AND LARRAIN, P. I. Evaluation of chemical and cultural control for noctuid larvae in Chilean artichokes produced for foreign markets, 115
MAH, J., LINZ, G. M. AND HANZEL, J. J. Relative effectivness of individual sunflower traits for reducing red-winged blackbird depredation, 359
MAHUKU, G. S. *see* LEVY, C., 148
MARTIN, A. P. *see* HENDERSON, I. F., 131
MARTIN, P. J. Comparison of the growth and yielding of clove trees under different ring weeding and fertilizer regimens, 415
MARTIN, T. D. *see* BLAIR, A. M., 289
MAUDE, R. B. *see* JORDAN, M. M., 367
MCCAFFERY, A. R. *see* TANZUBIL, P. B., 383
MCDONALD, D. *see* SHARMA, S. B., 453
MEAGHER JR., R. L. AND MEYER, J. R., Effects of ground cover management on certain abiotic and biotic interactions in peach orchard ecosystems, 65
MENASHEROV, M. *see* PASTER, N., 235
MEYER, J. R. *see* MEAGHER JR., R. L., 65
MOHAN, S. K. *see* LEITE JR, R. P., 3
MORA, M. *see* PASTER, N., 235
MPOFU, B. *see* CHIVINGE, O. A., 429

NACHMIAS, A. *see* ORDENTLICH, A., 363

O'MAKINWA, R. O. AND AKINYEMIJU, O. A. Control of *Euphorbia heterophylla*

L. in cowpea with herbicides and herbicide mixtures, 218
ORDENTLICH, A., NACHMIAS, A. AND CHET, I. Integrated control of *Verticillium dahliae* in potato by *Trichoderma harzianum* and captan, 363

PALMER, A. *see* SOPP, P. I., 177, 295
PANNELL, D. J. Model of wheat yield response to application of diclofop-methyl to control ryegrass (*Lolium rigidum*), 422
PAPAVIZAS, G. C. *see* LEWIS, J. A., 8
PARKER, K. A. *see* HENDERSON, I. F., 131
PARKIN, C. S. AND SIDDIQUI, H. A. Measurement of drop spectra from rotary cage aerial atomizers, 33
PASTER, N., CALDERON, M., MENASHEROV, M. AND MORA, M. Biogeneration of modified atmospheres in small storage containers using plant wastes, 235
PAYNE, P. A AND WILLIAMS, G. E. Hymexazol treatment of sugar-beet seed to control seedling disease caused by *Pythium* spp. and *Aphanomyces cochlioides*, 371
PEDIGO, L. P. *see* DAVIS, P. M., 387
PENROSE, L. J. Prolonged field persistence of resistance to benomyl in *Monilinia fructicola*, 190
PERKINS, T. J. *see* THORNTON, P. K., 337
PETTIT, T. R. *see* RUDGARD, S. A., 281
PRICE, D. *see* LINFIELD, C. A., 143
PRINGLE, K. L. *see* DRINKWATER, T. W., 128

RIEDER, P. *see* GOTSCH, N., 83
ROGNSTAD, A. *see* LUND-HØIE, K., 52, 59
ROMERO-MUÑOZ, F. *see* CASTEJON-MUÑOZ, M., 332
RUDGARD, S. A., PETTIT, T. R. AND HADLEY, P. Tenacity, biological activity and redistribution of copper fungicides on cocoa in controlled environments, 281

SAMOUCHA, Y. *see* COHEN, Y., 15
SANJRANI, W. *see* SUTHERLAND, J. A., 343
SCHNATHORST, W. C. *see* BRODY, A. K., 174
SEHGAL, V. K. AND UJAGIR, R. Effect of synthetic pyrethroids, neem extracts and other insecticides for the control of pod damage by *Helicoverpa armigera* (Hubner) on chickpea and pod damage-yield relationship at Pantnagar in northern India 29
SEUTIN, E. *see* VULSTEKE, G., 395
SHABI, E. *see* FREEMAN, S., 312
SHARMA, H. C. AND LOPEZ, V. F. Biology and population dynamics of sorghum head bugs (Hemiptera: Miridae), 164
SHARMA, S. B. AND MCDONALD, D. Global status of nematode problems of groundnut, pigeonpea, chickpea, sorghum, and pearl millet and suggestions for future work, 453
SIDDIQUI, H. A. *see* PARKIN, C. S., 33
SMITH, A. E. Potential allelopathic influence of certain pasture weeds, 410
SOPP, P. I. AND PALMER, A. Deposition patterns and biological effectiveness of spray deposits on pot plants applied by the Ulvafan and three prototype electrostatic sprayers, 295
SOPP, P. I., GILLESPIE, A. T. AND PALMER, A. Comparison of ultra-low-volume electrostatic and high-volume hydraulic application of *Verticillium lecanii* for aphid control in chrysanthemums, 177
SUETT, D. L. AND JUKES, A. A. Some factors influencing the accelerated degradation of mephosfolan in soils, 44
SUMMERS, R. W. AND HILLMAN, G. Scaring brent geese *Branta bernicla* from fields of winter wheat with tape, 459
SUTHERLAND, J. A., KING, W. J., DOBSON, H. M., INGRAM, W. R., ATTIQUE, M. R. AND SANJURANI, W. Effect of application volume and method on spray operator contamination by insecticide during cotton spraying, 343
SZTEJNBERG, A. *see* FREEMAN, S., 312

TAMAKI, G. *see* FOLWELL, R. J., 259
TANZUBIL, P. B. AND MCCAFFERY, A. R. Effects of azadirachtin and aqueous neem seed extracts on survival, growth and development of the African armyworm, *Spodoptera exempta*, 383
TATTERSFIELD, J. R. *see* LEVY, C., 148
THORNTON, P. K., FAWCETT, R. H., DENT, J. B. AND PERKINS, T. J. Spatial weed distribution and economic thresholds for weed control, 337

UJAGIR, R. *see* SEHGAL, V. K., 29
UKWUNGWU, M. N. Efficacy of granular isazofos against rice gall midge in Nigeria, 252

VAN EMDEN, H. F. *see* CLEAR HILL, B. H., 97
VAN DE BAAN, H. E., CROFT, B. A. AND BURTS, E. C. Resistance to the pyrethroid fenvalerate in pear psylla, *Psylla pyricola* Foerster (Homoptera: Psyllidae), in the Northwestern USA, 185
VELUSAMY, R. Relative susceptibility of high-yielding rice varieties to the thrips, *Stenchaetothrips biformis* (Bagnall) (Thysanoptera: Thripidae), 193
VULSTEKE, G. AND SEUTIN, E. Control of carrot fly (*Psila rosae* F.) in celeriac, 395

WALKER, A. *see* BLAIR, A. M., 289
WATT, A. D. *see* WRATTEN, S. D., 73
WELCH, S. J. *see* BLAIR, A.M., 289
WESTERN, N. M. *see* COOKE, B. K., 303
WICKS, T. J. Glashouse and field evaluation of fungicides for the control of *Septoria apiicola* on celery, 433
WILLIAMS, G. E. *see* PAYNE, P. A., 371
WRATTEN, S. D., WATT, A. D., CARTER, N. AND ENTWISTLE, J. C. Economic consequences of pesticide use for grain aphid control on winter wheat in 1984 in England, 73

YOUNG, A. G. Assessment of slug activity using bran-baited traps, 355

ŽĎÁRKOVÁ, E. AND HORÁK, E. Preventive biological control of stored food mites in empty stores using *Cheyletus eruditus* (Schrank), 378